


AD-A034 526


RHODE ISLAND UNIV KINGSTON AGRICULTURAL EXPERIMENT S--ETC F/G 6/18  
INFLUENCE OF EXTREMELY LOW FREQUENCY ELECTRIC AND MAGNETIC FIEL--ETC(U)  
DEC 76 W K DURFEE, C POLK, L T SMITH N00014-68-A-0215-0009  
CONTRIB-1729 NL

UNCLASSIFIED

| OF |

AD  
A034526





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS - 1963 - A

ADA 034526

Extremely Low Frequency Electric and Magnetic  
Fields in Domestic Birds

Final Report  
Phase II  
(Modulated Wave)

Departments  
of  
Animal Science  
and  
Electrical Engineering

University of Rhode Island  
Kingston, RI  
December, 1976

12  
B.S.

DDC  
RECEIVED  
JAN 18 1977  
RECEIVED

27

DISTRIBUTION STATEMENT A  
Approved for public release,  
Distribution Unlimited

⑥ Influence of Extremely Low Frequency  
Electric and Magnetic Fields Upon Growth  
Development and Behavior in Domestic Birds

⑨ Final Report - 15 Sep 71 - 30 Apr 76)

Phase II.

(Modulated Wave).

Submitted by

⑩ Wayne K. Durfee, Principal Investigator<sup>1</sup>

Charles Polk, Project Associate<sup>2</sup>

Lewis T. Smith, Project Associate<sup>1</sup>

Thomas J. Keefe, Research Assistant<sup>2</sup>

S. Muthukrishnan, Graduate Assistant<sup>2</sup>

- 
1. Department of Animal Science
  2. Department of Electrical Engineering

University of Rhode Island ✓  
Kingston, Rhode Island 02881

⑪ 15 Dec 76

⑫ 48p.

⑭ Contrib-1729

⑮

Supported in part (81 percent) by Naval Electronic Systems Command  
under Office of Naval Research Contract No. N00014-68-A-0215-0009  
cost-shared (19 percent) by University of Rhode Island

|                                 |                       |   |
|---------------------------------|-----------------------|---|
| DATE                            | 15 Dec 76             | ✓ |
| BY                              | Wayne K. Durfee       | ✓ |
| DISTRIBUTION/AVAILABILITY CODES |                       |   |
| REF.                            | APPL. ENG. OR SPECIAL |   |
| A                               |                       |   |

305550

6pg

Contribution number 1729, Rhode Island Agricultural Experiment Station.



## INTRODUCTION

The research herein reported was a continuation of that covered by Technical Report, Phase I (Continuous Wave) which was compiled March 1, 1975 (1).

The continuous wave studies were conducted with chick embryos, cultured chick embryo cells, immature chicks and adult chickens. Continuous wave frequencies of 45, 60 and 74 hertz were used; magnetic field intensities were 1, 5, 8 and 30 gauss; electric fields were 1, 10 and 3600 volts/meter. While chick embryo cell growth (in vitro) appeared to be inhibited by all but one ELF field studied, "no other responses (intact birds) were of sufficient magnitude or consistent enough to support the belief that they were due to the Elf fields tested."

The research in this report utilized the same special equipment and facilities which were designed for the earlier (CW) studies and which were easily adaptable for use with the modulated field frequencies in which the Navy was interested at the time the research was planned. In addition to the equipment used for the CW studies, specially designed equipment was supplied by the Navy to provide modulated signals intended to simulate the transmission from an ELF communication system.

## ABSTRACT

The domestic fowl was used in a series of experiments designed to allow evaluation of the influence of continuous exposure to modulated extremely low frequency (ELF) fields of low intensity upon 1) growth and development of the chick embryo; 2) early post-natal growth of the chick; 3) subsequent growth of the sexually immature bird; and 4) social interaction among sexually mature females.

The fields used were modulated between 72 and 80 hertz ( $76 \pm 4$  Hz). The magnetic fields were maintained at intensities of 1 or 8 gauss. The electric fields were maintained at either 10 V/m; 1 V/m; 10 V/m plus 60 Hz, 3.5 V/m; or 1 V/m plus 60 Hz, 3.5 V/m. Uniform fields at identical frequencies and amplitudes were provided for continuous exposure of embryos and chicks during the preincubation holding period, incubation and hatching periods and through the first 28 days of brooding.

Statistical analyses of the results of these studies revealed that neither the magnetic nor the electric fields tested had significant or consistent effects on the following:

1. Hatchability of fertile eggs.
2. Embryo survival during the most critical stages of development.
3. Early post-hatching growth and development.
4. Growth and development of the sexually immature bird.
5. Carbon dioxide production of the developing embryo.
6. Subsequent aggressiveness (as adults) of females exposed early in life.
7. Hematocrits of birds at 4-6 weeks or at 8 months of age.

The fields may have prolonged the incubation period slightly, but not enough to interfere with hatchability. More precise procedures would be necessary to better evaluate the effects of exposure to these fields upon the length of the incubation period and upon the growth of chicks which had been removed from field exposure soon after hatching.

The ELF fields used do not appear to have been detrimental to the survival or well-being of the exposed chicks.

## TECHNICAL DESCRIPTION OF THE WORK

The research described in this report was conducted in the facilities described earlier in Technical Report, Phase I (Continuous Wave) (1) and (2). These reports described the special equipment which was used to provide the uniform extremely low frequency (ELF) electric and magnetic fields at identical frequencies and amplitudes for continuous exposure of embryos and chicks throughout the preincubation holding, incubation, and hatching periods and through the first four weeks of brooding.

The earlier continuous wave work involved magnetic fields at frequencies of 45, 60 and 75 hertz and electric fields at 60 and 75 hertz. The magnetic fields were maintained at 1, 5, 8 or 30 gauss and the electric fields were maintained at 1, 10 and 3600 volts per meter. These ELF fields had no significant or consistent effects on: 1) hatchability of fertile eggs; 2) embryonic survival during the most critical stages of development; 3) early post-embryonic growth (to four weeks of age); and 4) learning and memory consolidation in the neo-nate chick. Growth and development to ten weeks of age was not affected by earlier (four-week) exposure. A 60 hertz, 5 gauss magnetic field had no effect on metabolic activity of chick embryos as determined by embryo growth rate and  $\text{CO}_2$  production.

Chick embryo cells grown in vitro appeared to be inhibited when incubated in 60 Hz magnetic fields at 1, 5 or 8 gauss; in 60 Hz electric fields of 1 or 10 V/m and in a 75 Hz electric field at 1 V/m. However, a 75 Hz electric field at 10 V/m appeared to accelerate cell growth. No explanation was offered for these seemingly opposite effects of the two 75 Hz fields. No other responses (intact birds) were of sufficient magnitude or consistent enough to support the belief that they were due to the ELF fields tested.



Upon completion of the continuous wave studies, it seemed desirable to gather additional data on the responses of the fowl to modulated ELF fields maintained at similar amplitudes to those used in the CW studies. Consequently, it was decided to use modulated frequencies close to 75 hertz. Existing special equipment for holding (preincubation) hatching eggs, incubation, hatching and rearing chicks to four weeks of age required modification only in electrical inputs and this added to the economic feasibility of these extended studies.

Four MSK generators were supplied by the Naval Electronic Systems Command from the IIT Research Institute to provide the desired signal modulation when properly connected to the ELF field simulators. The desired modulation parameters were " a pseudo random modulation pattern utilizing peak switching and a bit rate of 16. For these conditions and a center frequency of 76 Hz, the generator will switch between 72 and 80 Hz."<sup>1</sup> Subsequent references throughout this report will identify this modulated frequency as  $76 \pm 4$  hertz. The amplitudes of the modulated magnetic fields proposed for study were 1 and 8 gauss and those for the electric fields were 1 and 10 volts per meter. It was also planned to study the effects of the electric fields in the presence of continuous wave 60 hertz fields at 3.5 volts per meter.

#### OBJECTIVES

The objectives of the ELF modulated field studies were:

A. To evaluate the influence of continuous exposure (from egg pickup, through holding, incubation, and chick rearing to four weeks of age) to the following six fields:

---

<sup>1</sup>Communication from M. H. Benedick, Trip Report, URI, 7 February, 1975.



- 1)  $76 \pm 4$  Hz, 8 gauss
- 2)  $76 \pm 4$  Hz, 10 V/m
- 3)  $76 \pm 4$  Hz, 1 gauss
- 4)  $76 \pm 4$  Hz, 1 V/m
- 5)  $76 \pm 4$  Hz, 10 V/m plus 60 Hz CW, 3.5 V/m
- 6)  $76 \pm 4$  Hz, 1 V/m plus 60 Hz Cw, 3.5 V/m

B. To evaluate the influence of ELF fields 1), 2) and 5) above on the development of adult social behavior of hens continuously exposed during early development (prior to 4 weeks of age). (Field 5 was not used because the birds were immature until late January, 1976).

C. To evaluate the influence of ELF fields 1), 2) and 5) above upon carbon dioxide output of developing chick embryos.

#### MATERIALS AND METHODS

##### 1. Magnetic and Electric Field Simulators.

The ELF field simulators used in the earlier continuous wave studies were also used for these modulated frequency experiments. The incubator modifications and the hatcher-brooders used in both series (CW and modulated) can be seen in Figures 1-5. Detailed descriptions of the field simulators and the internal environments which prevailed in each have been presented earlier (1).

##### 2. The Egg Supply.

Hatching eggs from flock mated pens of White Plymouth Rocks (WPR) and White Leghorn (WL) fowl, maintained at the R. I. Agricultural Experiment Station poultry farm were used in these studies. White Plymouth Rocks were used in the trials which involved chick growth and subsequent adult be-




Figure 1. Exterior of Jamesway Model 252 incubators modified internally to accomodate ELF field simulators.




Figure 2. Interior view of Jamesway Model 252 incubator showing one electric field simulator in place at the top; a magnetic field simulator is at the center. Similar field simulators are in place in the lower part of the compartment.




Figure 3. Magnetic field simulator (1 of 2) for hatching eggs and brooding chicks to four weeks. One chick compartment and dropping tray are shown partly removed; the other compartment is accessible from the other end.




Figure 4. Electric field simulators used for hatching and brooding chicks through four weeks. Four plastic chick compartments are shown in place in one of the two heated E-field chambers. Each chamber is partitioned vertically to restrict field exposure to one side only.






Figure 5. Top view of one electric field chick compartment with furnishings in place showing; A, plastic chick compartment, 35" long, 8.5" wide; B, waterer, 2-quart; C, feeder; D, electric field probe (attached to feeder); E, plastic grid base (droppings pass through to plywood tray beneath).

havior because of the rapid growth rate customarily associated with chicks of this breed.

Leghorn eggs were used in the carbon dioxide studies because the smaller eggs from these birds could be used with less difficulty in the respiration chambers available for use with the infra-red CO<sub>2</sub> analyzer. (Each cylindrical respiration chamber had an inside diameter of 1.75 inches and was 8 inches long).

### 3. Coding of Experiments.

The Phase II Modulated Field studies were considered a continuation of the earlier CW experiments so a similar coding system was used for bird and data identification. The magnetic field trials were labeled M 2- and the electric field trials were labeled E 2-. The ELF field exposures used in Phase II are shown in Table 1 with the appropriate trial identification code, trial period dates, and brief notes about the nature of each trial.

### 4. Egg Handling and Body Weight.

All eggs collected daily from the breeder pens were placed in an egg holding room where they were maintained at approximately 60-70°F and 60-80% relative humidity until sufficient numbers were available for a complete setting. The holding period for each setting did not exceed ten days. The magnetic field experiments used 352 eggs and the electric field trials used 384 eggs for a complete set.

Within 24 hours after delivery to the holding room, each day's eggs were divided into equal numbers and stored in separate areas so that equal numbers of eggs of the same age would be set in the control or energized fields of the incubators. Eggs which were intended for ELF field exposure during incubation were continuously exposed to the same field during the preincubation holding period.



TABLE 1. SUMMARY OF MODULATED ELECTRIC AND MAGNETIC  
FIELD EXPOSURE STUDIES

| Trial<br>Code* | ELF Field<br>Exposure                        | Trial Period Dates |         |                |                 | Study                          |
|----------------|--|--------------------|---------|----------------|-----------------|--------------------------------|
|                |  | Set                | Hatched | Age<br>28 days | Age<br>10 weeks |                                |
| M 2-1          | 76 Hz $\pm$ 4, 8 gauss                       | 1/31/75            | 2/21    | 3/21           | 5/2             | Body Weight;<br>Adult Behavior |
| M 2-2          | 76 Hz $\pm$ 4, 1 gauss                       | 3/ 7/75            | 3/28    | 4/25           | 6/6             | Body Weight                    |
| E 2-1          | 76 Hz $\pm$ 4, 10 V/m                        | 2/17/75            | 3/10    | 4/7            | 5/19            | Body Weight;<br>Adult Behavior |
| E 2-2          | 76 Hz $\pm$ 4, 1 V/m                         | 3/24/75            | 4/14    | 5/12           | 6/23            | Body Weight                    |
| E 2-3          | 76 Hz $\pm$ 4, 10 V/m<br>+ 60 Hz CW, 3.5 V/m | 7/ 3/75            | 7/24    | 8/21           | 10/2            | Body Weight                    |
| E 2-4          | 76 Hz $\pm$ 4, 1 V/m<br>+ 60 Hz CW, 3.5 V/m  | 8/ 7/75            | 8/28    | 9/25           | 11/6            | Body Weight                    |
| M 2-3          | 76 Hz $\pm$ 4, 8 gauss                       | 5/11/75            | 6/1     |                |                 | CO <sub>2</sub>                |
| E 2-5          | 76 Hz $\pm$ 4, 10 V/m                        | 6/ 3/75            | 6/24    |                |                 | CO <sub>2</sub>                |
| E 2-6          | 76 Hz $\pm$ 4, 10 V/m<br>+ 60 Hz CW, 3.5 V/m | 6/12/75            | 6/27    |                |                 | CO <sub>2</sub>                |

\*M = magnetic field  
E = electric field

When enough eggs (approximately 400) had been collected for a complete setting, they were identified as to treatment and candled to allow removal of cracked or porous shelled eggs which would be unsuitable for hatching. The remaining eggs were then sorted in such a way that equal numbers of eggs of each age (days held before incubation) occurred in each simulator and in the same relative position within each simulator. This was accomplished by dividing each magnetic field cylinder into two 12-egg end zones and four 16-egg internal zones, and by subdividing each of the four electric field capacitors into six 16-egg zones.

Data used for comparisons between controls and ELF field-exposed embryos and chicks were collected from the eggs set in the internal zones. Chicks from the end zone eggs were subsequently reared under more "conventional" environmental conditions to provide an estimate of the effects of the special brooders (other than the ELF field effects) on early development of the chicks.

After 17 or 18 days of incubation (eggs set on Fridays were transferred on Mondays; eggs set on Mondays were transferred on Fridays), all eggs were removed from the incubators, candled to allow removal of infertile eggs and dead embryos, and all eggs with living embryos were transferred to hatcher-brooder compartments which were subdivided to allow continuation of the incubator setting zone designation through the hatching period.

Appropriate ELF field exposures, identical to those used during the preincubation and incubation periods, were continued during hatching and until the chicks reached four weeks of age.

Visual observations of the hatching compartments were made on the 21st day, starting at 496 hours (20 days, 16 hours) after setting, and

continuing at four-hour intervals until 512 hours (21 days, 8 hours). The chicks which had hatched at each observation time were identified by a series of toe-web marks and recorded to allow an estimation of the influence of the ELF fields on overall embryonic development.

On the 23rd day after setting, the chicks which had hatched were wing banded for individual identification, weighed, and the end zone chicks were moved to a starting battery. Unhatched eggs were placed under refrigeration, with those previously removed at transfer time, for later breakout to determine time of embryo death.

At seven days of age, all chicks were weighed individually and chick populations in the ELF field simulators were adjusted by removal, whenever possible, of chicks with odd-numbered bands. This plan should have resulted in the following distribution of chicks (a thermostat malfunction causing high mortality in E 2-1 and a miscount in M 2-2 resulted in smaller starting numbers in those trials):

#### Magnetic Field Simulators

##### Compartment

|   |         |           |
|---|---------|-----------|
| A | Exposed | 20 chicks |
| B | Control | 20 chicks |
| C | Control | 20 chicks |
| D | Exposed | 20 chicks |

#### "Conventional" Battery Compartment

|   |                                |
|---|--------------------------------|
| E | Excess from A and D, 20 chicks |
| F | Excess from B and C, 20 chicks |

## Electric Field Simulators

## Compartment

|                                 |                                    |
|---------------------------------|------------------------------------|
| A <sub>1</sub> , A <sub>2</sub> | Exposed, 10 chicks each; 20 chicks |
| B <sub>1</sub> , B <sub>2</sub> | Control, 10 chicks each; 20 chicks |
| C <sub>1</sub> , C <sub>2</sub> | Control, 10 chicks each; 20 chicks |
| D <sub>1</sub> , D <sub>2</sub> | Exposed, 10 chicks each; 20 chicks |

## "Conventional" Battery Compartments

|   |   |
|---|---|
| G | Excess from A <sub>1</sub> , A <sub>2</sub> , D <sub>1</sub> , D <sub>2</sub> ; 20 chicks |
| H | Excess from B <sub>1</sub> , B <sub>2</sub> , C <sub>1</sub> , C <sub>2</sub> ; 20 chicks |

At 28 days of age, after weighing, all remaining chicks were transferred from the field simulators or starting battery units to floor pens where all birds of each trial were reared together. The birds in trials M 2-1 and E 2-1 were reared to maturity for behavioral studies; all others were marketed after 10 weeks of age.

5. Carbon Dioxide Production of Embryos

Three experiments were conducted to evaluate the effects of one magnetic field and two electric fields (see Table 1) upon intact chick embryos.

The electronic gas analyzer (Beckman, Model IR 315) described earlier (1) was used to measure carbon dioxide output of intact chick embryos at 2-day intervals from day 6 through the 20th day of incubation.

Eggs from flock mated White Leghorns were used and for purposes of ELF field exposure were treated the same as those used for chick growth studies. Several handling differences were necessary, however, for these studies.

One incubator only was used for each study, and because the field simulators were located one above the other in each incubator (see Figure 2) the eggs were switched from one field simulator to another at two day intervals



after measuring  $\text{CO}_2$  output. This was intended to eliminate differences in  $\text{CO}_2$  output which may have resulted from slight temperature and other position differences within the incubator compartment. Electrical input connectors for each field allowed easy change of energized simulators when the eggs were changed. Continuous monitoring of each field, periodic recording of energized simulators, and color coding of exposed and control egg trays helped avoid switching errors.

At days, 6, 8, 10, 12, 14, 16, 18 and 20 equal numbers of exposed and control eggs were transferred hourly from the field simulators to a holding incubator (negligible ELF field) located in the room housing the gas analyzer. At approximately 15-minute intervals four eggs (two exposed and two control) were placed individually in the four heated analysis chambers. Two analysis chambers remained sealed throughout to provide continuous monitoring of instrument drift and warn of air leaks in the system. (Since  $\text{CO}_2$  content of the room air was high, any leaks allowed room air to blend with that from the analysis chambers and caused dramatic increases in recorded  $\text{CO}_2$ ). Figure 6 shows the general arrangement of the  $\text{CO}_2$  collection chambers and associated tubing. The chambers were enclosed in a compartment heated to approximately  $100^\circ\text{F}$  to minimize temperature fluctuations of the developing embryos which would have resulted from handling the eggs at room temperature ( $68\text{--}74^\circ\text{F}$ ). Except for the one-hour period required for  $\text{CO}_2$  analysis at 2-day intervals, all exposed eggs remained in the designated ELF field throughout each trial period.






Figure 6. Diagram showing arrangement of CO<sub>2</sub> collection chambers and associated tubing used with the gas analyzer.

## 6. Behavioral Studies

Adult females which had previously (until four weeks post-hatching) been exposed to an ELF magnetic field (M 2-1,  $76 \pm 4$  Hz, 8 gauss) or electric field (E 2-1,  $76 \pm 4$  Hz, 10 V/m) were used in peck order trials to determine whether or not the early field exposures had influenced their aggressiveness as adults. Four trials were conducted and each trial consisted of establishing a peck order among ten birds. Two trials were conducted among the females exposed as chicks to the magnetic field and two were conducted among those exposed as chicks to the electric field.

The pullets used were approximately 8-9 months of age and were in egg production when the trials were conducted. Each trial consisted of ten birds which had been placed in individual cages, to erase previous peck order hierarchies, three weeks prior to starting the test trials. Each trial was composed of ten birds; five were from control groups and were numbered 1-5, and five were from exposed groups and were numbered 6-10. After spending three weeks in cages without physical contact with one another, all birds in a trial were placed together in a 10-foot x 12-foot floor pen. Adequate food and water were available at all times. Daily observations of bird activities were made until a peck order had been established (usually within 3-4 days).

The data were analyzed by Chi-Square and Spearman's Rank Correlation methods (4).

## 7. Hematocrits

The hematocrit (packed cell volume expressed as percent of total blood) is low in young chickens and increases slightly in females and dramatically in males as they approach sexual maturity. Since circulating levels of testosterone increase hematocrits as well as aggressiveness in chickens, the hematocrit has been proposed as an indicator

of social aggressiveness in this species. Because hematocrit determinations are relatively simple to make, it was felt that this measure might have some potential value in these studies, particularly since some of the birds were to be reared to maturity. Therefore hematocrit determinations were made on samples of chicks which had been reared through four weeks of age in either a magnetic (M 2-1) or an electric (E 2-1) ELF field. Hematocrits were also determined on samples of the M 2-1 birds after they had reached sexual maturity (at approximately eight months of age).

The hematocrits were determined on small samples of blood drawn from a subcutaneous wing vein. (Several prominent veins are readily accessible as they cross or run parallel to the ulna, radius or humerus on the ventral side of the wing). The blood was collected in heparinized micro-hematocrit tubes, plugged with "Seal-Ease"<sup>1</sup> and centrifuged. Hematocrits were read with a "Spiro-Crit" tube reader.

## RESULTS AND DISCUSSION

The reader should refer to Table 1 for associating the assigned trial code numbers (M 2-1, E 2-1, etc.) with the ELF magnetic or electric field description for which they were used.

### 1. Hatchability.

The effects of the various ELF fields upon embryonic chick development were examined from the standpoints of hatchability of fertile eggs set (Table 2) embryo mortality by incubation period (Table 3) and hatching

---

<sup>1</sup>Micro-hematocrit tubes, "Seal-Ease", "Spiro-Crit" reader and micro-hematocrit centrifuge were products of Clay Adams, Division of Becton, Dickinson and Company, Parsippany, NJ 07054.

TABLE 2. COMPARISON OF HATCHABILITY OF FERTILE EGGS SET BETWEEN CONTROLS AND EGGS EXPOSED TO VARIOUS MODULATED ELECTRIC AND MAGNETIC FIELDS DURING THE HOLDING AND INCUBATION PERIODS

| Trial Code | Treatment Group | No. Eggs Set | Fertile Eggs | Chicks Hatched <sup>1</sup><br>No. | %    | chi-Square Value (DF=1) | P     |
|------------|-----------------|--------------|--------------|------------------------------------|------|-------------------------|-------|
| M 2-1      | Exposed         | 128          | 108          | 67                                 | 62.0 | 3.27                    | > .05 |
|            | Control         | 128          | 106          | 78                                 | 73.6 |                         |       |
| M 2-2      | Exposed         | 128          | 120          | 74                                 | 61.7 | 1.83                    | > .10 |
|            | Control         | 128          | 126          | 88                                 | 69.8 |                         |       |
| E 2-1      | Exposed         | 128          | 123          | 78                                 | 63.4 | 0.95                    | > .30 |
|            | Control         | 128          | 117          | 67                                 | 57.3 |                         |       |
| E 2-2      | Exposed         | 128          | 117          | 80                                 | 68.4 | .29                     | > .50 |
|            | Control         | 128          | 123          | 88                                 | 71.5 |                         |       |
| E 2-3      | Exposed         | 128          | 105          | 71                                 | 67.6 | ----- <sup>2</sup>      |       |
|            | Control         | 128          | ---          | 77                                 | ---  |                         |       |
| E 2-4      | Exposed         | 128          | 110          | 77                                 | 70.0 | 1.63                    | > .20 |
|            | Control         | 128          | 99           | 61                                 | 61.6 |                         |       |

<sup>1</sup>Number and Percent of Fertile Eggs Hatched.

<sup>2</sup>Control eggs were discarded without checking for embryo mortality.



TABLE 3. COMPARISON OF MORTALITY BY INCUBATION PERIODS BETWEEN CONTROLS AND EMBRYOS EXPOSED TO VARIOUS MODULATED ELECTRIC AND MAGNETIC FIELDS

| Trial Code | Treatment Group    | Eggs Set   | Chicks Hatched | Early Deaths <sup>1</sup> | Mortality by Incubation Period (Days) |        |          | Chi-square Value (DF=3) | P     |
|------------|--------------------|------------|----------------|---------------------------|---------------------------------------|--------|----------|-------------------------|-------|
|            |                    |            |                |                           | 1-7                                   | 7-14   | 14-21    |                         |       |
| M 2-1      | Exposed<br>Control | 128<br>128 | 67<br>78       | 20<br>22                  | 19<br>13                              | 1<br>1 | 21<br>14 | 1.56                    | > .50 |
| M 2-2      | Exposed<br>Control | 128<br>128 | 74<br>88       | 7<br>2                    | 35<br>23                              | 2<br>0 | 9<br>15  | 7.28                    | > .05 |
| E 2-1      | Exposed<br>Control | 128<br>128 | 78<br>67       | 5<br>11                   | 21<br>33                              | 1<br>2 | 23<br>15 | 5.90                    | > .10 |
| E 2-2      | Exposed<br>Control | 128<br>128 | 80<br>88       | 11<br>5                   | 20<br>16                              | 1<br>0 | 16<br>19 | 3.25                    | > .30 |
| E 2-3      | Exposed<br>Control | 128<br>128 | 71<br>77       | 23                        | 16<br>--                              | 3<br>- | 15<br>-- | <sup>2</sup><br>-----   |       |
| E 2-4      | Exposed<br>Control | 128<br>128 | 77<br>61       | 18<br>29                  | 19<br>25                              | 0<br>1 | 14<br>12 | 2.42                    | > .30 |

<sup>1</sup>Includes infertile and mortality prior to incubation.

<sup>2</sup>Control eggs were discarded without checking for embryo mortality.



time in terms of hours after setting (Table 4, 5 and 6). In examining these data it should be kept in mind that the eggs used in four trials (M 2-1, M 2-2, E 2-1 and E 2-2) were gathered, held, and incubated during cold weather; those used in two trials (E 2-3 and E 2-4) were gathered, held and incubated in July and August. In addition, the rate of egg production of the breeders necessitated holding the eggs longer in most cases (up to two days) than is the customary commercial practice (usually 5-6 days) and these factors probably accounted for the low hatchability experienced in these studies. However, the identical practices followed with the control and exposed eggs (other than the experimental variables under study) made these data suitable for analysis.

Hatchability of fertile eggs set (Table 2) appeared to be unaffected by the ELF fields used and mortality by incubation period (days 1-7, 7-14, and 14-21) were similarly unaffected (Table 3). Hatchability differences between exposed and control eggs approached significance in only one trial (M 2-1) and differences in mortality by incubation period approached significance in only one trial (M 2-2).

Since embryo survival did not appear to follow a pattern in favor of either the exposed or control condition, however, it seems unlikely that these differences could be attributed to the ELF field exposures.

In conducting the earlier studies (Phase I, Continuous Wave) the observation had been made, without supporting data, that the eggs which had been incubated in one magnetic field coil seemed to hatch earlier than either the control eggs or those from the other energized coil. Therefore, an attempt was made to determine whether or not differences existed in hatching time between controls and eggs exposed

TABLE 4. COMPARISON OF HATCHING TIME BETWEEN CONTROLS AND EMBRYOS EXPOSED TO VARIOUS MODULATED ELECTRIC AND MAGNETIC FIELDS

| Trial Code <sup>1</sup> | Treatment Group | Eggs Set   | Chicks hatched by Hatching Time |               |          | Total Chicks Hatched | Chi-Square Value | P DF=2     |
|-------------------------|-----------------|------------|---------------------------------|---------------|----------|----------------------|------------------|------------|
|                         |                 |            | Early                           | Inter-mediate | Late     |                      |                  |            |
| M 2-1                   | Exposed Control | 128<br>128 | 20<br>41                        | 15<br>17      | 32<br>20 | 67<br>78             | 9.34             | .01 >.0005 |
| M 2-2                   | Exposed Control | 128<br>128 | 15<br>23                        | 16<br>25      | 43<br>40 | 74<br>88             | 2.58             | .30 >.25   |
| E 2-1                   | Exposed Control | 128<br>128 | 21<br>17                        | 22<br>23      | 35<br>27 | 78<br>67             | 0.65             | .75 >.70   |
| E 2-2                   | Exposed Control | 128<br>128 | 16<br>28                        | 30<br>39      | 34<br>21 | 80<br>88             | 7.15             | .05 >.025  |
| E 2-4                   | Exposed Control | 128<br>128 | 54<br>50                        | 22<br>9       | 1<br>2   | 77<br>61             | 4.14             | .20 >.10   |

<sup>1</sup>See Table 1 for Trial Code identification.

<sup>2</sup>Hatching Time (21 days = 504 hours)

Early: < 500 hours after setting.  
Intermediate: 500-508 hours after setting.  
Late: > 508 hours after setting.

TABLE 5. COMPARISON OF HATCHING TIME BETWEEN ALL EGGS INCUBATED IN TWO TOP SIMULATORS AND THOSE INCUBATED IN TWO BOTTOM SIMULATORS

| Trial Code <sup>1</sup> | Incubator Position | Eggs Set | Chicks hatched by Hatching Time <sup>2</sup> |               |      | Total Chicks Hatched | Chi-Square Value | P DF=2     |
|-------------------------|--------------------|----------|--|---------------|------|----------------------|------------------|------------|
|                         |                    |          | Early  | Inter-mediate | Late |                      |                  |            |
| M 2-1                   | Top                | 128      | 30   | 16            | 16   | 62                   | 4.77             | .10 > .05  |
|                         | Bottom             | 128      | 31   | 16            | 36   | 83                   |                  |            |
| M 2-2                   | Top                | 128      | 23   | 16            | 37   | 76                   | 4.04             | .20 > .10  |
|                         | Bottom             | 128      | 15   | 25            | 46   | 86                   |                  |            |
| E 2-1                   | Top                | 128      | 26   | 30            | 18   | 74                   | 21.00            | .001 >     |
|                         | Bottom             | 128      | 12   | 15            | 44   | 71                   |                  |            |
| E 2-2                   | Top                | 128      | 33   | 37            | 18   | 88                   | 17.58            | .001 >     |
|                         | Bottom             | 128      | 11   | 32            | 37   | 80                   |                  |            |
| E 2-4                   | Top                | 128      | 62   | 10            | 0    | 72                   | 10.51            | .01 > .005 |
|                         | Bottom             | 128      | 42   | 21            | 3    | 66                   |                  |            |

<sup>1</sup> See Table 1 for Trial Code identification.

<sup>2</sup> Hatching time (21 days = 504 hours).

Early: < 500 hours after setting.

Intermediate: 500-508 hours after setting.

Late: > 508 hours after setting.

TABLE 6.  
COMPARISON OF HATCHING TIME BETWEEN CONTROLS AND EMBRYOS EXPOSED TO  
VARIOUS MODULATED ELECTRIC AND MAGNETIC FIELDS WHEN HATCHING TIME IS BASED ON  
APPROXIMATE MEAN HATCHING TIME OF CONTROLS

| Trial<br>Code <sup>1</sup> | Treatment<br>Group | Eggs<br>Set | Chicks Hatched by                                     |                |               |                             | Chi-Square<br>Value | P<br>DF=1 |
|----------------------------|--------------------|-------------|---|----------------|---------------|-----------------------------|---------------------|-----------|
|                            |                    |             | Control Mean<br>Hatching Time<br>(Hours) <sup>2</sup> | Hatching Time  |               | Number<br>Chicks<br>Hatched |                     |           |
|                            |                    |             |   | Early<br>No. % | Late<br>No. % |                             |                     |           |
| M 2-1                      | Exposed<br>Control | 128         | 500   | 20 29.9        | 47 70.1       | 67                          | 7.63                | .01 >.005 |
|                            |                    | 128         |   | 41 52.6        | 37 47.4       | 78                          |                     |           |
| M 2-2                      | Exposed<br>Control | 128         | 504   | 23 31.1        | 51 68.9       | 74                          | 3.50                | .10 >.05  |
|                            |                    | 128         |   | 40 45.5        | 48 54.5       | 88                          |                     |           |
| E 2-1                      | Exposed<br>Control | 128         | 504   | 29 37.2        | 49 62.8       | 78                          | .32                 | .70 >.50  |
|                            |                    | 128         |   | 28 41.8        | 39 58.2       | 67                          |                     |           |
| E 2-2                      | Exposed<br>Control | 128         | 504   | 30 37.5        | 50 62.5       | 80                          | 3.69                | .10 >.05  |
|                            |                    | 128         |   | 46 52.3        | 42 47.7       | 88                          |                     |           |
| E 2-4                      | Exposed<br>Control | 128         | 496   | 34 44.2        | 43 55.8       | 77                          | 3.71                | .10 >.05  |
|                            |                    | 128         |   | 37 60.7        | 24 39.3       | 61                          |                     |           |

<sup>1</sup>See Table 1 for Trial Code identification.

<sup>2</sup>Hours after setting at which 42-61% of the control chicks had hatched; time used to distinguish early from late hatched chicks.



to the various fields. The chicks which hatched were recorded at four-hour intervals during hatching and the analyses of the data collected appear in Tables 4-6. The numbers of chicks which hatched early, intermediate and late are presented, together with the Chi-Square analysis of the data.

The differences between the control and exposed eggs were great enough in two trials (M 2-1 and E 2-2) to encourage further observations of the data. Therefore, the data were compared by position (top simulator vs. bottom) in the incubators with exposed and control data pooled. This analysis indicated a position effect (Table 5) and prompted the analysis shown in Table 6 which shows a comparison between exposed and control embryos when early and late hatch designations depend upon approximately 50 percent hatch of the controls. These data are not precise because of the difficulty of measuring time at 50 percent hatch with the procedure used (observations at 4-hour intervals), however they show a trend toward a prolongation of the hatch which may have been due to slight retardation of development during incubation. In order to evaluate more precisely the influence of the ELF fields on this response, greater care would be necessary to eliminate position and temperature differences during incubation. While these factors did not appear to affect eventual hatchability of the fertile eggs, it is well known that slight temperature differences during the full 21-day incubation period can have a dramatic effect (retardation or acceleration by several hours) on the time of hatch.

The observations at hatching time of E 2-3 were inadequate for analysis. The differences in hatching time between the control and exposed chicks in E 2-1 were not significant, but differences between

TABLE 7 . COMPARISONS BETWEEN BIRD BODY WEIGHTS OF CONTROLS AND CHICKS  
EXPOSED TO A MAGNETIC FIELD OF  $76 \pm 4$  Hz, 8 gauss  
(M 2-1) THROUGH 28 DAYS OF AGE

| Bird Age (Days) | Treatment                    | Number of Birds | Bird Body Weight (grams) |                    |          | Degrees of Freedom | Computed t Value |
|-----------------|------------------------------|-----------------|--------------------------|--------------------|----------|--------------------|------------------|
|                 |                              |                 | Mean                     | Standard Deviation | Variance |                    |                  |
| 2               | Exposed Control              | 40              | 37.1                     | 4.17               | 17       | 78                 | 0.7259           |
|                 |                              | 40              | 37.7                     | 3.18               | 10       |                    |                  |
| 7               | Exposed Control              | 39              | 48.3                     | 10.25              | 105      | 77                 | 0.5388           |
|                 |                              | 40              | 47.2                     | 6.73               | 45       |                    |                  |
| 14              | Exposed Control              | 37              | 90.5                     | 21.85              | 478      | 74                 | 0.4904           |
|                 |                              | 39              | 88.4                     | 15.81              | 250      |                    |                  |
| 21              | Exposed Control              | 39              | 150.3                    | 38.78              | 1504     | 77                 | 0.0314           |
|                 |                              | 40              | 150.5                    | 30.09              | 906      |                    |                  |
| 28              | Exposed Control              | 40              | 237.5                    | 41.48              | 1721     | 78                 | 0.1386           |
|                 |                              | 40              | 238.8                    | 46.14              | 2129     |                    |                  |
| 70              | Exposed <sup>2</sup> Control | 39              | 1470.1                   | 315.37             | 99461    | 76                 | 0.7909           |
|                 |                              | 39              | 1413.8                   | 318.75             | 101599   |                    |                  |

<sup>1</sup> Critical t value at .05 level of significance and 75 degrees of freedom = 1.995.

<sup>2</sup> No field exposure after 28 days of age.

TABLE 8. COMPARISONS BETWEEN BIRD BODY WEIGHTS OF CONTROLS AND CHICKS  
EXPOSED TO A MAGNETIC FIELD OF  $76 \pm 4$  Hz, 1 gauss  
(M 2-2) THROUGH 28 DAYS OF AGE

| Bird<br>Age<br>(Days) | Treatment            | Number<br>of<br>Birds | Bird Body Weight (grams) |                       |          | Degrees<br>of<br>Freedom | Computed<br>t Value |
|-----------------------|----------------------|-----------------------|--------------------------|-----------------------|----------|--------------------------|---------------------|
|                       |                      |                       | Mean                     | Standard<br>Deviation | Variance |                          |                     |
| 2                     | Exposed              | 40                    | 40.0                     | 3.52                  | 12       | 77                       | 1.8495              |
|                       | Control              | 39                    | 38.5                     | 3.91                  | 15       |                          |                     |
| 7                     | Exposed              | 40                    | 53.9                     | 4.07                  | 16       | 77                       | 2.5895*             |
|                       | Control              | 39                    | 50.9                     | 6.07                  | 37       |                          |                     |
| 14                    | Exposed              | 40                    | 99.4                     | 10.69                 | 114      | 77                       | 1.0056              |
|                       | Control              | 39                    | 96.8                     | 12.68                 | 161      |                          |                     |
| 21                    | Exposed              | 40                    | 146.8                    | 17.59                 | 310      | 77                       | 0.9287              |
|                       | Control              | 39                    | 150.7                    | 19.66                 | 387      |                          |                     |
| 28                    | Exposed              | 40                    | 253.4                    | 29.09                 | 846      | 77                       | 0.4458              |
|                       | Control              | 39                    | 256.4                    | 31.30                 | 980      |                          |                     |
| 70                    | Exposed <sup>2</sup> | 34                    | 1326.2                   | 585.77                | 343131   | 70                       | 0.0387              |
|                       | Control              | 38                    | 1322.0                   | 311.39                | 96965    |                          |                     |

<sup>1</sup> Critical t value at .05 level of significance and 75 degrees of freedom = 1.995.

\* Significant at the .02 level.

<sup>2</sup> No field exposure after 28 days of age.

TABLE 9. COMPARISONS BETWEEN BIRD BODY WEIGHTS OF CONTROLS AND CHICKS  
EXPOSED TO AN ELECTRIC FIELD OF  $76 \pm 4$  Hz, 10 V/m  
(E 2-1) THROUGH 28 DAYS OF AGE

| Bird Age<br>(Days) | Treatment | Number<br>of<br>Birds | Bird Body Weight (grams) |                       |          | Degrees<br>of<br>Freedom | Computed<br>t Value <sup>1</sup> |
|--------------------|-----------|-----------------------|--------------------------|-----------------------|----------|--------------------------|----------------------------------|
|                    |           |                       | Mean                     | Standard<br>Deviation | Variance |                          |                                  |
| 2                  | Exposed   | 35                    | 36.2                     | 12.9                  | 166      | 64                       | 0.0488                           |
|                    | Control   | 31                    | 36.4                     | 13.7                  | 188      |                          |                                  |
| 7                  | Exposed   | 31                    | 58.8                     | 31.1                  | 970      | 70                       | 0.6917                           |
|                    | Control   | 41                    | 55.2                     | 12.0                  | 145      |                          |                                  |
| 14                 | Exposed   | 37                    | 122.5                    | 34.7                  | 1200     | 76                       | 0.6097                           |
|                    | Control   | 41                    | 118.5                    | 22.9                  | 526      |                          |                                  |
| 21                 | Exposed   | 39                    | 229.6                    | 37.6                  | 1416     | 76                       | 0.8857                           |
|                    | Control   | 39                    | 218.9                    | 66.4                  | 4411     |                          |                                  |
| 28                 | Exposed   | 39                    | 387.2                    | 58.6                  | 3431     | 78                       | 1.7351                           |
|                    | Control   | 41                    | 361.9                    | 71.9                  | 5176     |                          |                                  |
| 84                 | Exposed   | 38                    | 2418.2                   | 474.8                 | 225471   | 71                       | 0.4166                           |
|                    | Control   | 35                    | 2343.2                   | 1009.4                | 1018971  |                          |                                  |

<sup>1</sup>Critical t value at .05 level of significance and 75 degrees of freedom = 1.995.

<sup>2</sup>No field exposure after 28 days of age.



TABLE 10. COMPARISONS BETWEEN BIRD BODY WEIGHTS OF CONTROLS AND CHICKS  
EXPOSED TO AN ELECTRIC FIELD OF  $76 \pm 4$  Hz, 1 V/m  
(E 2-2) THROUGH 28 DAYS OF AGE

| Bird Age (Days) | Treatment            | Number of Birds | Bird Body Weight (grams) |                    |          | Degrees of Freedom | Computed t Value |
|-----------------|----------------------|-----------------|--------------------------|--------------------|----------|--------------------|------------------|
|                 |                      |                 | Mean                     | Standard Deviation | Variance |                    |                  |
| 2               | Exposed              | 40              | 36.3                     | 3.3                | 11       | 78                 | 2.0975*          |
|                 | Control              | 40              | 37.8                     | 3.4                | 12       |                    |                  |
| 7               | Exposed              | 40              | 50.5                     | 7.6                | 57       | 78                 | 0.8722           |
|                 | Control              | 40              | 51.8                     | 6.3                | 39       |                    |                  |
| 14              | Exposed              | 40              | 101.4                    | 18.3               | 336      | 78                 | 0.6191           |
|                 | Control              | 40              | 103.6                    | 14.9               | 222      |                    |                  |
| 21              | Exposed              | 40              | 198.8                    | 33.5               | 1122     | 78                 | 0.1168           |
|                 | Control              | 40              | 197.9                    | 32.0               | 1021     |                    |                  |
| 28              | Exposed              | 40              | 318.5                    | 47.0               | 2206     | 78                 | 0.9927           |
|                 | Control              | 40              | 328.6                    | 44.6               | 1986     |                    |                  |
| 70              | Exposed <sup>2</sup> | 20              | 1386.3                   | 1149.3             | 1320933  | 40                 | 0.0807           |
|                 | Control              | 22              | 1411.4                   | 876.5              | 768175   |                    |                  |

<sup>1</sup>Critical t value at .05 level of significance and 75 degrees of freedom = 1.995.

\* Significant at .05 level.

<sup>2</sup>No field exposure after 28 days of age.

TABLE 11. COMPARISONS BETWEEN BIRD BODY WEIGHTS OF CONTROLS AND CHICKS EXPOSED TO AN ELECTRIC FIELD OF  $76 \pm 4$  Hz, 10 V/m plus 60 Hz CW, 3.5 V/m (E 2-3) THROUGH 28 DAYS OF AGE

| Bird Age (Days) | Treatment            | Number of Birds | Bird Body Weights (grams) |                    |          | Degrees of Freedom | Computed t Value <sup>1</sup> |
|-----------------|----------------------|-----------------|---------------------------|--------------------|----------|--------------------|-------------------------------|
|                 |                      |                 | Mean                      | Standard Deviation | Variance |                    |                               |
| 2               | Exposed              | 40              | 38.8                      | 3.0                | 9        | 78                 | 0.0703                        |
|                 | Control              | 40              | 38.7                      | 3.4                | 11       |                    |                               |
| 7               | Exposed              | 40              | 63.6                      | 8.0                | 64       | 78                 | 0.8035                        |
|                 | Control              | 40              | 62.2                      | 7.1                | 50       |                    |                               |
| 14              | Exposed              | 40              | 110.1                     | 16.0               | 256      | 77                 | 2.2679*                       |
|                 | Control              | 39              | 100.2                     | 22.6               | 509      |                    |                               |
| 21              | Exposed              | 39              | 187.7                     | 51.1               | 2616     | 74                 | 1.2110                        |
|                 | Control              | 37              | 171.9                     | 62.8               | 3948     |                    |                               |
| 28              | Exposed              | 39              | 313.1                     | 85.2               | 7253     | 73                 | 1.1491                        |
|                 | Control              | 36              | 287.3                     | 110.6              | 12234    |                    |                               |
| 70              | Exposed <sup>2</sup> | 31              | 1413.1                    | 714.1              | 509915   | 54                 | 0.0685                        |
|                 | Control              | 25              | 1399.0                    | 840.3              | 706166   |                    |                               |

<sup>1</sup> Critical t value at .05 level of significance and 75 degrees of freedom = 1.995.

\* Significant at .05 level

<sup>2</sup> No field exposure after 28 days of age.

TABLE 12. COMPARISONS BETWEEN BIRD BODY WEIGHTS OF CONTROLS AND CHICKS  
EXPOSED TO AN ELECTRIC FIELD OF  $76 \pm 4$  Hz, 1 V/m  
plus 60 Hz CW, 3.5 V/m (E 2-4) THROUGH 28 DAYS OF AGE

| Bird Age (Days) | Treatment            | Number of Birds | Bird Body Weight (grams) |                    |          | Degrees of Freedom | Computed t Value |
|-----------------|----------------------|-----------------|--------------------------|--------------------|----------|--------------------|------------------|
|                 |                      |                 | Mean                     | Standard Deviation | Variance |                    |                  |
| 2               | Exposed              | 40              | 37.0                     | 3.4                | 12       | 78                 | 1.8476           |
|                 | Control              | 40              | 35.9                     | 3.1                | 10       |                    |                  |
| 7               | Exposed              | 40              | 67.8                     | 6.8                | 46       | 78                 | 0.0961           |
|                 | Control              | 40              | 67.6                     | 7.2                | 52       |                    |                  |
| 14              | Exposed              | 40              | 145.2                    | 15.7               | 247      | 77                 | 0.0791           |
|                 | Control              | 39              | 144.8                    | 28.7               | 821      |                    |                  |
| 21              | Exposed              | 40              | 249.3                    | 31.8               | 1010     | 77                 | 0.1313           |
|                 | Control              | 39              | 248.1                    | 48.5               | 2351     |                    |                  |
| 28              | Exposed              | 40              | 398.8                    | 49.3               | 2430     | 77                 | 0.1917           |
|                 | Control              | 39              | 401.6                    | 78.4               | 6148     |                    |                  |
| 70              | Exposed <sup>2</sup> | 34              | 1304.4                   | 487.1              | 237245   | 67                 | 0.1957           |
|                 | Control              | 35              | 1282.1                   | 464.7              | 215957   |                    |                  |

<sup>1</sup>Critical t value at .05 level of significance, 75 degrees of freedom.

<sup>2</sup>No field exposure after 28 days of age.

controls and exposed chicks in the remaining four trials indicate that the ELF fields retarded embryo development slightly.

## 2. Bird Body Weight

After all viable chicks had hatched (approximately 23 days after setting), they were wing banded for individual identification and weighed. They were again weighed individually at 7, 14, 21, 28 and 70 days after hatching (exception: the birds in trial E 2-1 were weighed at 12 weeks rather than at ten weeks of age). The body weight data and their analyses are shown in Tables 7-12 for each growth trial. Since individual data were available, the analyses made use of this fact, avoiding the need to use the means derived from replicated groups.

The variation in bird numbers whose body weights were available for analysis at the various periods was due to several factors. In several instances, lost wing bands were replaced without body weight determinations at the appropriate time but in such cases body weight was taken at the next regular time. Chicks which died within the first two weeks were considered abnormal and their body weights were not analyzed. In the case of trial E 2-1, a malfunction of one of the hatcher-brooder temperature controls caused overheating of the unit and loss of most of the chicks at day 8. Since the surplus chicks had been removed only one day earlier, the lost chicks were replaced by returning surplus chicks to the experimental chambers from the battery brooder maintained in the same room. The exposed chicks in this trial were therefore outside the experimental ELF field for approximately 20 hours.

In a few instances, an occasional chick was overlooked on "weigh day", presumed dead, then discovered on the succeeding weigh day. In such cases only the available data as recorded were analyzed, but bird



TABLE 13. BODY WEIGHTS OF BIRDS EXPOSED TO VARIOUS ELF MAGNETIC OR ELECTRIC FIELDS THROUGH SEVEN DAYS OF AGE AND REARED IN BATTERY COMPARTMENTS THROUGH 28 DAYS OF AGE

| Trial Code | Early Treatment Group <sup>1</sup> | Number Chicks @ |         | Body Weight (grams) |         |                      |
|------------|------------------------------------|-----------------|---------|---------------------|---------|----------------------|
|            |                                    | 2 days          | 70 days | 2 days              | 28 days | 70 days <sup>2</sup> |
| M 2-1      | Exposed                            | 17              | 15      | 36.8                | 256.2   | 1550.3               |
|            | Control                            | 18              | 18      | 32.9                | 277.8   | 1478.3               |
| M 2-2      | Exposed                            | 20              | 20      | 39.0                | 374.4   | 1507.5               |
|            | Control                            | 20              | 19      | 37.8                | 335.2   | 1405.0               |
| E 2-1      | Exposed                            | 17              | 16      | 38.0                | 367.5   | 2549.1               |
|            | Control                            | 20              | 19      | 38.4                | 365.7   | 2444.2               |
| E 2-2      | Exposed                            | 20              | 8       | 38.7                | 381.9   | 1500.0               |
|            | Control                            | 20              | 16      | 37.7                | 353.3   | 1448.1               |
| E 2-3      | Exposed                            | 20              | 18      | 38.7                | 318.8   | 1418.9               |
|            | Control                            | 20              | 13      | 38.5                | 319.0   | 1315.5               |
| E 2-4      | Exposed                            | 20              | 18      | 37.1                | 390.8   | 1326.7               |
|            | Control                            | 20              | 13      | 36.5                | 403.6   | 1285.8               |

<sup>1</sup>No birds were exposed to ELF fields after 7 days of age.

<sup>2</sup>Body weights of E 2-1 birds were at 84 days of age.

numbers in the tables appear to fluctuate. In several cases, a sharp decrease in bird numbers between 28 and 70 days is obvious (especially E 2-2 and E 2-3). This was due to the mortality caused mostly by coccidiosis which is particularly troublesome in flocks started on grills which prevent ready access to their droppings during early development. Such birds develop little resistance to coccidia and when placed on litter often experience heavy mortality. Proper preventive medication of such birds is difficult.

Body weight differences between exposed and control chicks were great enough to be significant at the .05 level in only three instances:

in trial M 2-2, Exposed > Control at 7 days;

in trial E 2-2, Control > Exposed at 2 days;

in trial E 2-3, Exposed > Control at 14 days;

These data do not support a hypothesis that bird body weight prior to ten weeks of age was influenced by the ELF electric or magnetic fields studied.

Body weights of surplus chicks removed at seven days and reared in battery compartments are shown in Table 13. Birds from the exposed groups were reared in compartments separate from the ones from the control groups. Body weights at 48 hours, 28 days and 70 days are shown. When these body weight data are compared with the ones in Tables 7-12 they indicate that a population density of 20 chicks per magnetic field simulator compartment and 10 chicks per electric field simulator compartment allowed growth responses comparable to those observed in the battery compartments (20 chicks per unit).

At 70 days of age (84 days for E 2-1) the chicks which originated from the energized groups (until 7 days post-hatching) had greater body

weights than comparable control chicks. Since exposure to the ELF fields was terminated at 7 days of age, and because other exposed (through 28 days of age) chicks did not show consistently similar differences from their controls, it is difficult to explain the apparently greater growth of the previously exposed, battery reared chicks. Approximately equal numbers of previously exposed and control chicks were used in each battery compartment to eliminate position effects (chicks were identified by band number).

### 3. Carbon Dioxide Production of Embryos.

The mean carbon dioxide production of chick embryos from day 6 through day 20 were summarized in Tables 14-16 for trials M 2-3, E 2-5, and E 2-6. The data were gathered from individual embryos whose identity was maintained throughout the trial period. The tables include analyses of the differences in carbon dioxide production between control embryos and those continuously exposed to a modulated magnetic field of 8 gauss (M 2-3) a modulated electric field of 10 V/m (E 2-5) or a modulated electric field of 10 V/m plus 60 Hz CW, 3.5 V/m (E 2-6). The data show repeated measurements on the same embryos and therefore if the ELF fields had influenced metabolic rate of carbon dioxide production, it is reasonable to assume that the effects would have been cumulative or at least would have shown a trend. The data do not indicate any such trend.

In trial M 2-3, the exposed embryos exceeded the controls slightly throughout; the difference was significant only on day 18.

In trial E 2-5, the exposed embryos produced  $\text{CO}_2$  at a more rapid rate than the controls on days 6, 8, 10 and 16; on days 12, 14, 18 and 20 the controls produced more  $\text{CO}_2$ .

TABLE 14. COMPARISON BETWEEN MEAN CO<sub>2</sub> PRODUCTION (ml CO<sub>2</sub>/egg/hr) OF CONTROLS  
AND EMBRYOS EXPOSED TO A MODULATED MAGNETIC FIELD

76 ± 4 Hz, 8 gauss

| Day | Treatment | Number of<br>Embryos | Mean   | Standard<br>Deviation | Variance | Degrees of<br>Freedom | Computed<br>t value |
|-----|-----------|----------------------|--------|-----------------------|----------|-----------------------|---------------------|
| 6   | Exposed   | 19                   | 0.892  | 0.124                 | 0.0154   | 39                    | 1.7001              |
|     | Control   | 22                   | 0.826  | 0.127                 | 0.0161   |                       |                     |
| 8   | Exposed   | 10                   | 1.658  | 0.253                 | 0.064    | 20                    | 0.9884              |
|     | Control   | 12                   | 1.545  | 0.291                 | 0.085    |                       |                     |
| 10  | Exposed   | 19                   | 2.258  | 0.394                 | 0.155    | 38                    | 1.7337              |
|     | Control   | 21                   | 2.059  | 0.341                 | 0.116    |                       |                     |
| 12  | Exposed   | 18                   | 4.660  | 0.856                 | 0.733    | 37                    | 1.3409              |
|     | Control   | 21                   | 4.335  | 0.673                 | 0.453    |                       |                     |
| 14  | Exposed   | 19                   | 9.863  | 2.330                 | 5.428    | 39                    | 0.539               |
|     | Control   | 22                   | 9.539  | 1.516                 | 2.298    |                       |                     |
| 16  | Exposed   | 19                   | 15.335 | 1.579                 | 2.493    | 39                    | 0.7627              |
|     | Control   | 22                   | 14.758 | 3.002                 | 9.134    |                       |                     |
| 18  | Exposed   | 19                   | 21.783 | 2.252                 | 5.071    | 39                    | 3.0936*             |
|     | Control   | 22                   | 19.891 | 1.695                 | 2.872    |                       |                     |
| 20  | Exposed   | 18                   | 21.083 | 2.479                 | 6.144    | 36                    | 0.1804              |
|     | Control   | 20                   | 20.911 | 3.380                 | 11.421   |                       |                     |

<sup>1</sup>Critical t value at 5% level of significance, 20 degrees of freedom = 2.086  
Critical t value at 5% level of significance, 40 degrees of freedom = 2.021.

\* Significant at .005 level.



TABLE 15. COMPARISON OF MEAN CO<sub>2</sub> PRODUCTION (ml CO<sub>2</sub>/egg/hr) OF CONTROLS  
AND EMBRYOS EXPOSED TO MODULATED ELECTRIC FIELDS

76 ± 4 Hz, 10 V/m

| Day | Treatment          | Number of Embryos | Mean             | Standard Deviation | Variance         | Degrees of Freedom | Computed t value <sup>1</sup> |
|-----|--------------------|-------------------|------------------|--------------------|------------------|--------------------|-------------------------------|
| 6   | Exposed<br>Control | 43<br>49          | 0.747<br>0.712   | 0.218<br>0.144     | 0.0474<br>0.0209 | 90                 | 0.9216                        |
| 8   | Exposed<br>Control | 43<br>49          | 1.256<br>1.151   | 0.320<br>0.140     | 0.103<br>0.0196  | 90                 | 2.0811*                       |
| 10  | Exposed<br>Control | 44<br>48          | 2.172<br>2.081   | 0.396<br>0.344     | 0.157<br>0.119   | 90                 | 1.183                         |
| 12  | Exposed<br>Control | 43<br>49          | 4.187<br>4.227   | 0.638<br>0.561     | 0.408<br>0.314   | 90                 | 0.3216                        |
| 14  | Exposed<br>Control | 43<br>49          | 9.961<br>10.104  | 1.318<br>1.205     | 1.736<br>1.453   | 90                 | 0.5464                        |
| 16  | Exposed<br>Control | 43<br>49          | 17.599<br>17.236 | 1.593<br>1.691     | 2.537<br>2.859   | 90                 | 1.0618                        |
| 18  | Exposed<br>Control | 43<br>49          | 20.514<br>20.872 | 2.253<br>1.766     | 5.077<br>3.118   | 90                 | 0.8569                        |
| 20  | Exposed<br>Control | 42<br>47          | 21.837<br>22.921 | 3.173<br>2.820     | 10.065<br>7.950  | 87                 | 1.7156                        |

<sup>1</sup>Critical t value at 5% level of significance, 60 degrees of freedom = 2.000.

\* Significant at .05 level.

TABLE 16. COMPARISON OF MEAN CO<sub>2</sub> PRODUCTION (ml CO<sub>2</sub>/egg/hr) OF CONTROLS AND EMBRYOS EXPOSED TO A MODULATED ELECTRIC FIELD

76 ± 4Hz, 10 V/m, plus 60 Hz CW, 3.5 V/m

| Day | Treatment       | Number of Embryos | Mean             | Standard Deviation | Variance         | Degrees of Freedom | Computed t value <sup>1</sup> |
|-----|-----------------|-------------------|------------------|--------------------|------------------|--------------------|-------------------------------|
| 6   | Exposed Control | 45<br>37          | 1.059<br>0.992   | 0.321<br>0.154     | 0.103<br>0.0236  | 80                 | 1.1666                        |
| 8   | Exposed Control | 44<br>38          | 1.379<br>1.353   | 0.259<br>0.204     | 0.067<br>0.0418  | 80                 | 0.5014                        |
| 10  | Exposed Control | 42<br>38          | 2.709<br>2.742   | 0.313<br>0.268     | 0.098<br>0.072   | 78                 | 0.5063                        |
| 12  | Exposed Control | 44<br>37          | 5.962<br>5.856   | 0.689<br>0.769     | 0.475<br>0.591   | 79                 | 0.6588                        |
| 14  | Exposed Control | 44<br>37          | 12.139<br>12.662 | 1.113<br>1.156     | 1.239<br>1.337   | 79                 | 2.0833*                       |
| 16  | Exposed Control | 44<br>37          | 18.077<br>18.199 | 2.179<br>2.137     | 4.746<br>4.565   | 79                 | 0.2548                        |
| 18  | Exposed Control | 44<br>36          | 22.473<br>23.040 | 3.459<br>3.118     | 11.965<br>9.742  | 78                 | 0.7662                        |
| 20  | Exposed Control | 44<br>37          | 24.323<br>25.178 | 4.635<br>4.754     | 21.481<br>22.604 | 79                 | 0.8227                        |

<sup>1</sup> Critical t value for 5% level of significance, 60 degrees of freedom = 2.000.

\* Significant at .05 level.

In trial E 2-6, the exposed embryos produced  $\text{CO}_2$  at a higher rate than the controls on days 6, 8 and 12; on days 10, 14, 16, 18 and 20 they produced less  $\text{CO}_2$ .

The differences in  $\text{CO}_2$  production were less than significant (.05 level) on most days measured.

These data indicate that the ELF fields used did not alter metabolic rate of the chick embryos enough to be measured as a significant change in carbon dioxide production rate. While these data do not support the evidence cited earlier that ELF fields may have slowed embryonic development slightly, it seems reasonable to assume that embryonic development could have been slowed enough to retard hatching time by a few hours without significantly reducing  $\text{CO}_2$  production (considering individual embryo variation and the limits of precision imposed by the measuring procedures).

#### 4. Behavioral Studies.

Adult female White Plymouth Rock pullets which had been reared together, then isolated by housing in individual cages, were identified by their wing bands according to their earlier (through 28 days post-hatching) ELF magnetic or electric field exposure and were provided with readily visible wing badges numbered 1-10. Within each trial, birds numbered 1-5 were from control groups and those numbered 6-10 were from previously exposed (M 2-1 or E 2-1) groups. Frequent observations of the birds after recombining in floor pens by assigned numbers (1-10) revealed the establishment of new peck orders within three to four days. Table 17 shows the peck orders established in the four trials conducted. The birds were ranked by aggressiveness; the most aggressive was placed at the top and the least aggressive (most submissive) was placed at the bottom.

TABLE 17. COMPARISONS OF AGGRESSIVENESS BETWEEN ADULT PULLETS  
EXPOSED AS CHICKS (THROUGH 28 DAYS OF AGE) TO A  
MAGNETIC OR ELECTRIC ELF FIELD AND THEIR CONTROLS

| Rank of Aggressiveness by Bird Number <sup>1</sup> |  |          |   |          |
|--|--|----------|---|----------|
| Order of<br>Aggressiveness                         | Electric Field<br>E 2-1<br>76 $\pm$ 4 Hz, 10 V/m |          | Magnetic Field<br>M 2-1<br>76 $\pm$ 4 Hz, 8 gauss |          |
|  | Trial I  | Trial II | Trial III   | Trial IV |
|  |  |          |   |          |
| Most   | 6  | 9        | 8   | 6        |
|  | 1  | 1        | 9   | 10       |
|  | 9  | 2        | 1   | 9        |
|  | 2  | 6        | 4   | 8        |
|  | 3  | 8        | 10  | 4        |
|  | 8  | 4        | 6   | 1        |
|  | 4  | 5        | 3   | 3        |
|  | 5  | 10       | 7   | 2        |
|  | 7  | 7        | 2   | 7        |
| Least  | 10   | 3*       | 5   | 5        |

<sup>1</sup>In each trial, 1-5 were from control groups;  
6-10 were from previously exposed groups.

\*Bird No. 3 died on day 2 of the trial.



Chi square and Spearman's Rank Correlation analyses were used to test the significance of the results.

With the exception of Trial IV, the orders constituted a random order; Trial IV approached non-randomness. Combining all tests and testing for agreement among tests revealed "no agreement". This indicated that previous exposure to the ELF field tested did not influence peck order.

No correlation was found between peck order and body weight or between peck order and hematocrit at 4 or 6 weeks of age.

#### 5. Hematocrits.

The hematocrits determined on randomly selected birds from the groups previously exposed to either an Elf magnetic field (M 2-1,  $76 \pm 4$  Hz, 8 gauss) or to an ELF electric field (E 2-1,  $76 \pm 4$  Hz, 10 V/m) appear in Table 18. Determinations were made on the M 2-1 birds at 6 weeks and at 8 months of age; those on the E 2-1 birds were made at 4 weeks of age. No significant differences between the hematocrits of the previously exposed and control birds were found.

### CONCLUSIONS

The data gathered from a series of experimental trials in which chick embryos and neo-nate chicks were continuously exposed to modulated ELF magnetic or electric fields indicate that the fields investigated did not cause consistent or significant changes in the following:

1. Hatchability of fertile eggs.
2. Embryo survival during the most critical stages of development.
3. Early post-hatching growth and development.

TABLE 18. HEMATOCRITS OF IMMATURE AND ADULT CHICKENS EXPOSED TO AN ELF MAGNETIC  
OR ELECTRIC FIELD FOR FOUR WEEKS AFTER HATCHING

|              | Immature <sup>1</sup>           |         |                                |         | Adult <sup>2</sup>              |         |
|--------------|---------------------------------|---------|--------------------------------|---------|---------------------------------|---------|
|              | M 2-1                           |         | E 2-1                          |         | M 2-1                           |         |
|              | Exposed<br>76 ± 4 Hz<br>8 gauss | Control | Exposed<br>76 ± 4 Hz<br>10 V/m | Control | Exposed<br>76 ± 4 Hz<br>8 gauss | Control |
| Chick Number | 10                              | 10      | 9                              | 9       | 11                              | 11      |
| Hematocrit   |                                 |         |                                |         |                                 |         |
| Mean         | 33.3                            | 33.3    | 32.3                           | 31.7    | 36.6                            | 34.8    |
| s.d.         | 2.03                            |         | 2.03                           |         | 3.81                            |         |
| s.e.         | .91                             |         | .96                            |         | 1.62                            |         |
| t            | 0.0                             |         | .69                            |         | 1.07                            |         |
| D.F.         | 18                              |         | 16                             |         | 20                              |         |
| P            | >.9                             |         | .5-.6                          |         | .25-.30                         |         |

<sup>1</sup>M 2-1 chicks were 6 weeks old  
E 2-1 chicks were 4 weeks old

<sup>2</sup>Birds were approximately 8 months old.

4. Growth and development of the sexually immature bird.
5. Carbon dioxide production of the developing embryo.
6. Subsequent aggressiveness (as adults) of females exposed early in life.
7. Hematocrits of birds at 4-6 weeks or at 8 months of age.

The fields may have prolonged the incubation period slightly, but not enough to interfere with hatchability. More precise procedures would be necessary to better evaluate the effects of exposure to these fields upon the length of the incubation period and upon the growth of chicks which had been removed from field exposure soon after hatching.

The ELF fields used do not appear to have been detrimental to the survival or well-being of the exposed chicks.

## REFERENCES

1. Durfee, W. K., P. W. Chang, C. Polk, L. T. Smith, V. J. Yates, P. R. Plante, S. Muthukrishnan, and H. Chen, 1975. Influence of extremely low frequency electric and magnetic fields upon growth, development and behavior in domestic birds. Technical Report, Phase I (Continuous Wave). Entry Number B-15, Compilation of Navy Sponsored ELF Biomedical and Ecological Research Reports, Volume II. Naval Medical Research and Development Command, Department of the Navy, National Naval Medical Center, Bethesda, Md. 20014, February, 1975.
2. Durfee, W. K., P. R. Plante, P. Martin, S. Muthukrishnan, and C. Polk, 1975. Exposure of domestic fowl to ELF electric and magnetic fields. Paper B10a-5. Proceedings, 1975 Annual Meeting, USNC/URSI-IEEE. Univ. of Colorado, Boulder, Colo., Oct. 20-23. (In Press).
3. Hall, Jeffrey A., 1972. The Effects of Diethylstilbestrol, Testosterone Propionate, and Progesterone on the Hematocrit in Gallus Domesticus. M.S. Thesis, Univ. of R.I.
4. Ostle, B., 1963. Statistics in Research, 2nd ed. Iowa State University Press, Ames, Iowa.
5. Plante, Paul R., 1974. Evaluation of the Effect of Extremely Low Frequency Electric and Magnetic Fields upon Growth, Development, and Memory in Gallus Domesticus. M.S. Thesis, Univ. of R.I.



| REPORT DOCUMENTATION PAGE  |                       | READ INSTRUCTIONS<br>BEFORE COMPLETING FORM   |
|--|-----------------------|---|
| 1. REPORT NUMBER   | 2. GOVT ACCESSION NO. | 3. RECIPIENT'S CATALOG NUMBER   |
| 4. TITLE (and Subtitle)<br><b>Influence of Extremely Low Frequency Electric and Magnetic Fields Upon Growth, Development, and Behavior in Domestic Birds</b>   |                       | 5. TYPE OF REPORT & PERIOD COVERED<br><b>Final Report - Phase 2</b><br><b>Sept. 15, 1971-April 30, 1976</b> |
|  |                       | 6. PERFORMING ORG. REPORT NUMBER  |
| 7. AUTHOR(s)<br><b>Wayne K. Durfee, Principal Investigator</b><br><b>Charles Polk, Project Associate</b><br><b>Lewis T. Smith, Project Associate</b>   |                       | 8. CONTRACT OR GRANT NUMBER(s)<br><b>N00014-68-A-0215-0009</b> ✓  |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS<br><b>Department of Animal Science</b><br><b>214 Woodward Hall, University of Rhode Island</b><br><b>Kingston, Rhode Island 02881</b>  |                       | 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS<br><br><b>NR 201- 089</b>                       |
| 11. CONTROLLING OFFICE NAME AND ADDRESS<br><br><b>Naval Electronic Systems Command</b>   |                       | 12. REPORT DATE<br><b>December 15, 1976</b>   |
|  |                       | 13. NUMBER OF PAGES   |
| 14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)<br><b>Office of Naval Research</b><br><b>Arlington, Virginia 22217</b>   |                       | 15. SECURITY CLASS. (of this report)<br><b>Unclassified</b>   |
|  |                       | 15a. DECLASSIFICATION/DOWNGRADING SCHEDULE  |
| 16. DISTRIBUTION STATEMENT (of this Report)<br><br><b>Approved for public release; distribution unlimited.</b>   |                       |   |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)   |                       |   |
| 18. SUPPLEMENTARY NOTES  |                       |   |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number)<br><br><b>Extremely Low Frequency (ELF) Fields      Embryo Development</b><br><b>Electric Fields                                      Chick Growth</b><br><b>Magnetic Fields                                     Bird Behavior</b><br><b>Modulation</b>   |                       |   |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number)<br><br>✓ <b>Chicks and embryos of the domestic fowl were continuously exposed to ELF magnetic or electric fields modulated between 72 and 80 hertz. The magnetic fields were maintained at an intensity of 1 or 8 gauss and the electric fields were maintained at either 10 V/m; 1 V/m; 10 V/m plus 60 Hz, 3.5 V/m; or 1 V/m plus 60 Hz, 3.5 V/m. Uniform fields at identical frequencies and amplitudes were provided for continuous exposure of embryos</b><br><div style="text-align: right;">(cont on p 47) ✓</div> |                       |   |

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

6 cont  
and chicks and during the preincubation holding period, incubation and hatching periods and through the first 28 days of brooding.

Statistical analyses of the <sup>study</sup> results of these studies revealed that neither the magnetic nor the electric fields tested had significant or consistent effects on the following:

1. Hatchability of fertile eggs;
2. Embryo survival during the most critical stages of development;
3. Early post-hatching growth and development;
4. Growth and development of the sexually immature bird;
5. Carbon dioxide production of the developing embryo;
6. Subsequent aggressiveness (as adults) of females exposed early in life; or
7. Hematocrits of birds at 4-6 weeks or at 8 months of age.

The fields may have prolonged the incubation period slightly, but not enough to interfere with hatchability. More precise procedures would be necessary to better evaluate the effects of exposure to these fields upon the length of the incubation period and upon the growth of chicks which had been removed from field exposure soon after hatching.

The ELF fields used do not appear to have been detrimental to the survival or well-being of the exposed chicks.

